HIPPPO EDUCATION
















Objectives



Apply clinically relevant milestones to the approach to pediatric exams/procedures



Highlight key aspects to the approach, examination and immobilization of pediatric patients





• Appear calm and approachable

Model positive behavior with the guardian

Break the ICE





Age Related Considerations



Age	Milestone	Action
Infant (0-2 years old)	Stranger anxiety, easily startled, easily distracted	Warm hands, distractions, lean into parent interaction



Age	Milestone	Action
Toddler (2-3 years old)	Stubborn, minimal language, short attention span	Model calm behavior, short direct language, distraction





Age	Milestone	Action
Preschool (3-6 years old)	Improved language, emotionally reactive, social	Explain/involve without overwhelming





Age	Milestone	Action
School age (6-12 years old)	Body awareness, good at negotiation, cognitively	Direct language, involve patient, limit negotiations,



Preparation

- Set expectations
 - Discuss plan with parents
 - Define clear roles with your team
- Equipment
 - Have a plan and a back up plan
 - Gather materials beforehand
 - Keep sharps out of sight!

Distraction

- Tablet/Cell phone
- Toys
- Games
- Music
- Parents
- Guided imagery
- Child life specialist

Extra Analgesia

- Topicals
 - Creams/Gels
 - Freeze spray
- Oral medications
- Injectables






Analgesia	Time of onset
Lidocaine	1-3 minutes
Bupivacaine	5-10 minutes
EMLA	60 minutes
LET	15-30 minutes
Ethyl chloride	<30 seconds

Decreases pain, improves cooperation, and may avoid the use of needles

PMID: 31386316



Examination Techniques



Infant: Swaddle



Newborn/infant: swaddle





Infant > school age: side sitting








Older infant: back to chest



Toddler: chest to chest



Immobilization



Superhero Burrito Cape

PMID: 21296532







LESS IS MORE

TAKE A BREAK/DEBRIEFEND ON A POSITIVE NOTEBEFORE DISCHARGE

Bedside Magic

HIPPPO EDUCATION

The Floating Cup




Coin Magic



The Magic Wand

Special Considerations

- Patient selection
 - Read the room
- Prop selection
 - Sterility
- Presentation
 - Make the patient the star of the performance
 - Consider leaving the patient with a souvenir
 - Disposal of props



Practice Makes Perfect !

References

Brown, Julie and Klein, Eileen. The "Superhero Cape Burrito": A Simple and comfortable method of short-term procedural restraint. *The Journal of Emergency Medicine*. 2011, Vol 41 Issue 1 P74-76. Accessed November 11, 2023. <u>link</u>

Children's Hospital of Philadelphia (CHOP). *Comfort positioning during procedures*.Published July 10, 2017. Accessed November 11, 2023 Link

Darleu, Charlene. *How to Approach Pediatric Patients*. American Association of Critical Care Nurses. Published May 4, 2023. Accessed November 11, 2023 Link

Uspal N, Black KD, Cico SJ. Pediatric pain management in the emergency department. Pediatr Emerg Med Pract. 2019 Aug;16(8):1-24. Epub 2019 Aug 2. PMID: 31339255. Link

HIPPPO EDUCATION